

RKF IMPACTS SURVIVAL



7-12%

Lower risk of death associated with urine volume^{1,2,3}



6-36%

Lower risk of death associated with GFR^{1,4}

PRESERVING RKF



OVERHYDRATION

does not preserve RKF⁹ and can lead to left ventricular hypertrophy & higher risk of death^{5,6}



PHARMACOLOGY

using ACE, ARB, loop diuretics may be helpful⁷⁻¹⁰

www.AdvancingDialysis.org

1. Jansen MAM, Hart AAM, Korevaar JC, Dekker FW, Boeschoten EW, Krediet RT. Predictors of the rate of decline of residual renal function in incident dialysis patients. *Kidney Int.* 2002;62(3):1046-1053. doi:10.1046/j.1523-1755.2002.00505.x
2. Rumpsfeld M, McDonald SP, Johnson DW. Peritoneal Small Solute Clearance is Nonlinearly Related to Patient Survival in the Australian and New Zealand Peritoneal Dialysis Patient Populations. *Perit Dial Int J Int Soc Perit Dial.* 2009;29(6):637-646. doi:10.1177/089686080902900609
3. Korevaar, Johanna C., Boeschoten, Elisabeth W., Dekker, Friedo W., Krediet, Raymond T. for TNSG. What Have We Learned about PD from Recent Major Clinical Trials? *Perit Dial Int J Int Soc Perit Dial.* 2007;27(1):10-10. doi:10.1177/089686080702700103
4. Chen CH, Perl J, Teitelbaum I. Prescribing high-quality peritoneal dialysis: The role of preserving residual kidney function. *Perit Dial Int.* 2020;40(3):274-281. doi:10.1177/0896860819893821
5. Shu Y, Liu J, Zeng X, et al. The effect of overhydration on mortality and technique failure among peritoneal dialysis patients: A systematic review and meta-analysis. *Blood Purif.* 2018;46(4):350-358. doi:10.1159/000492148
6. Perl J, Davies SJ, Lambie M, et al. The peritoneal dialysis outcomes and practice patterns study (Pdopps): Unifying efforts to inform practice and improve global outcomes in peritoneal dialysis. *Perit Dial Int.* 2016;36(3):297-307. doi:10.3747/pdi.2014.00288
7. Suzuki H, Kanno Y, Sugahara S, Okada H, Nakamoto H. Effects of an angiotensin II receptor blocker, valsartan, on residual renal function in patients on CAPD. *Am J Kidney Dis.* 2004;43(6):1056-1064. doi:10.1053/j.ajkd.2004.01.019
8. Li PKT, Chow KM, Wong TYH, Leung CB, Szeto CC. Effects of an Angiotensin-Converting Enzyme Inhibitor on Residual Renal Function in Patients Receiving Peritoneal Dialysis: A Randomized, Controlled Study. *Ann Intern Med.* 2003;139(2). doi:10.7326/0003-4819-139-2-200307150-00010
9. Lameire N, Van Biesen W. Importance of blood pressure and volume control in peritoneal dialysis patients. *Perit Dial Int.* 2001;21(2):206-211. doi:10.1177/089686080102100219
10. Medcalf JF, Harris KPG, Walls J. Role of diuretics in the preservation of residual renal function in patients on continuous ambulatory peritoneal dialysis. *Kidney Int.* 2001;59(3):1128-1133. doi:10.1046/j.1523-1755.2001.0590031128.x