

CARDIOVASCULAR DISEASE




41+%

of all dialysis patient deaths
are cardiovascular-related¹



LVH

is an important predictor of
cardiovascular mortality and morbidity²



10% = 28%

decrease
in LV mass

lower risk
for cardiovascular
related death³

BOTTOM LINE?



17%

lower risk of cardiovascular death with
intensive as compared to conventional hemodialysis⁴

www.AdvancingDialysis.org

1. Saran R, Li Y, Robinson B, et al. US Renal Data System 2014 Annual Data Report: Epidemiology of Kidney Disease in the United States. Am J Kidney Dis. 2015;66(1 Suppl 1):Svii, S1-305. doi:10.1053/j.ajkd.2015.05.001.

2. Luthje L, Andreas S. Obstructive sleep apnea and coronary artery disease. Sleep Med Rev. 2008;12(1):19-31. doi:10.1016/j.smr.2007.08.002.

3. London, G.M., Pannier, B., Guerin, A.P. et al, Alterations of left ventricular hypertrophy in and survival of patients receiving hemodialysis: follow-up of an interventional study. J Am Soc Nephrol. 2001;12:2759-2767.

4. Weinhandl ED, Liu J, Gilbertson DT, Arneson TJ, Collins AJ. Survival in daily home hemodialysis and matched thrice-weekly in-center hemodialysis patients. JASN. 2012;23(5):895-904. doi:10.1681/ASN.2011080761